

## Parallel Implementation of the Self-Consistent-Charge Density-Functional-Based Tight-Binding

Ch. Köhler, M. Haugk, Z. Hajnal, A. Blumenau, A. Sieck, Th.  
Frauenheim

published in

*Modern Methods and Algorithms of Quantum Chemistry*,  
J. Grotendorst (Ed.), John von Neumann Institute for Computing,  
Jülich, NIC Series, Vol. 2, ISBN 3-00-005746-3, p. 21, 2000.

© 2000 by John von Neumann Institute for Computing

Permission to make digital or hard copies of portions of this work for personal or classroom use is granted provided that the copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. To copy otherwise requires prior specific permission by the publisher mentioned above.

<http://www.fz-juelich.de/nic-series/>

# Parallel Implementation of the Self-Consistent-Charge Density-Functional-Based Tight-Binding

Christof Köhler, Michael Haugk, Zoltan Hajnal, Alexander  
Blumenau, Alexander Sieck, Thomas Frauenheim

University of Paderborn, Department of Physics,  
33095 Paderborn, Germany

The density-functional-based tight-binding method with charge-selfconsistency (SCC-DFTB) has been applied to a variety of problems in the fields of solids, surfaces, inorganic clusters and even biological molecules. This fast and efficient method has proved to be very accurate, results are usually in very good agreement with fully self-consistent DFT results.

The method has been implemented for parallel computers based on MPI and Scalapack and was tested on various hardware including the T3E and PC-based distributed-memory computers. We present details about the performance and scaling behaviour and demonstrate applications of the parallel code.